

BRACHYTHERAPY

This Model Policy addresses coverage for Brachytherapy.

DESCRIPTION

Radiation oncology consists of two primary treatment modalities: external beam radiation therapy (EBRT) and brachytherapy. Brachytherapy is a type of radiation therapy that utilizes natural or manufactured radioactive isotopes or radionuclides that are temporarily or permanently implanted to treat malignancies or certain benign conditions and derives a physical advantage based upon the inverse square law of physics. Brachytherapy is accomplished by implanting a sealed radioactive source directly in or adjacent to the tumor or treatment site. The basic clinical brachytherapy formats are: superficial, interstitial, intracavitary or intraluminal therapy. "Electronic" brachytherapy, which utilizes miniature X-ray sources operating at low-kilovoltage energies to approximate dose distributions achieved with radionuclide brachytherapy for selected cases, is an emerging treatment modality and not subject to this Model Policy. Brachytherapy may use a solid radioactive source, such as a 'seed' or liquid colloid isotopes and may be either temporary or permanent. Further, brachytherapy may be called high-dose-rate or low-dose-rate depending on the dose delivery per hour.

Brachytherapy may be used independently as the sole treatment or as an adjunctive treatment in combination with external beam therapy and other modalities such as surgery or chemotherapy.

Brachytherapy may be performed concomitantly with surgical resection or in conjunction with procedures such as endoscopy or angioplasty, which are required to achieve access to the site of the disease. There are two distinct phases to the brachytherapy process: the insertion of radioelements through catheters, needles or other special devices called brachytherapy applicators, and the loading of the radioactive material (the active or therapeutic agent) into the conduits or directly into tissue.

The typical requirements of brachytherapy may involve:

- Treatment planning (77263)
Brachytherapy treatment planning is routinely designated complex (CPT code 77263) because it requires complex treatment volume design, dose levels near normal tissue tolerance, analysis or special tests, complex fractionation, or delivery concurrent with other therapeutic modalities or treatment of previously irradiated tissues.
- Dosimetry (77300)
Brachytherapy requires certain calculations to be made throughout the course of treatment. Each basic dosimetry calculation may be submitted when performed for brachytherapy treatment.
- Treatment management
The management of brachytherapy includes the overall management of the patient during the course of radiation therapy. It includes review of the treatment record to assure that the therapy is proceeding according to the radiation plan and dose prescription, medical evaluation of the patient during the course of treatment, making the necessary adjustments to the applicator during the course of treatment, and any other modifications or clinical interventions needed to assure safe and effective treatment during the clinical course of therapy.
- Special treatment procedure (77470)
The delivery of brachytherapy often requires special arrangements with the operating room, radiation safety team and/or inpatient ward, coordination with other specialists, preparation and provision of the applicators and related equipment, scheduling and integration of required physics support, and acquisition and preparation of the radiation sources. Brachytherapy is often delivered in conjunction with external radiation, chemotherapy or surgery. Integration of these processes makes brachytherapy a special treatment procedure.

¹ ASTRO model policies were developed to efficiently communicate what ASTRO believes to be correct coverage policies for radiation oncology services. The ASTRO Model Policies do not serve as clinical guidelines and they are subject to periodic review and revision without notice. The ASTRO Model Policies may be reproduced and distributed, without modification, for noncommercial purposes.

- Simulation (77280-77290)
For brachytherapy, simulation may require the use of imaging examinations of the implanted sources or applicator(s) containing dummy (i.e., non-radioactive) sources. These films of the implanted sources are used to verify quality of applicator positioning, develop isodose curves and other dosimetry, and may be billed separately, when appropriate.
- Isodose planning (77316-77318), (77295)
Brachytherapy requires an isodose plan. This plan determines the dose at each implanted source and throughout the treatment volume and doses to surrounding normal tissue. CPT Codes 77316, 77317, and 77318 include basic dosimetry, so 77300 cannot be billed with these codes. CPT code 77295 may be billed as part of the brachytherapy process when the needed parameters are included (i.e., 3-D volume reconstruction with DVH for target and normal tissues, etc.). The NCCI procedure to procedure edit for CPT Codes 77300 and 77295 was deleted in October 2016.
- Handling and Loading of Radioelement (77790)
Where brachytherapy techniques require the manual loading of an isotope (LDR), the supervision, loading and handling of the isotope may be separately reported. (Do not report in conjunction with 77778).
- Source Application/Placement (77750, 77761-77763, 77767-77772, 77778 and 77789)
Selection and placement of after-loading applicators and the loading and unloading of radioactive sources may be performed by the radiation oncologist alone or in collaboration with another physician.
- Applicator placement
The choice of applicators and the actual placement of the after-loading device may be performed by the radiation oncologist alone or in collaboration with another physician (e.g., gynecologist, urologist, pulmonologist).
- Treatment devices (77332-77334)
Treatment devices may include the use of certain templates, molds or other apparatus that may be required for specific clinical circumstances. These may be covered as treatment devices.
- Medical Physics Services (77336, 77370, 77331)
CPT code 77336 is a “weekly code;” however, for radiation therapy treatment that is not administered in five weekly fractions (such as brachytherapy) or for a course of radiation therapy consisting of one or two fractions, code 77336 may be reported.

CPT code 77370 may be justified for the complex interrelationships of electron and photon ports and complex dosimetric considerations in brachytherapy, including high-dose-rate remote afterloader applications, intravascular brachytherapy treatments and interstitial radioactive seed implantation.

CPT code 77331 is Special dosimetry (e.g., TLD, microdosimetry) when prescribed by a physician. In some instances, measurement of the delivered radiation dose may be used to guide and determine the dose to selected positions within or around the implant treatment volume.

CPT/HCPCS CODES:

Many of the CPT codes in the radiation oncology section can be used for both external beam radiation therapy (EBRT) and brachytherapy, while others are specific to one modality or the other. The CPT codes below, with the exception of 77261-77263, 77280-77295, 77336 and 77370 are those CPT codes specific only to brachytherapy. CPT codes 77300 (Basic Dosimetry), 77332-77334 (Treatment devices, designs and construction), 77336 (Continuing medical physics consultation, including assessment of treatment parameters, quality assurance of dose delivery and review of patient treatment documentation in support of the radiation oncologist, reported per week of therapy), 77370 (Special medical radiation physics consultation) and 77470 (Special treatment procedure) are also relevant to brachytherapy.

77261 Therapeutic radiology treatment planning; simple

77262 ; intermediate

77263 ; complex

77280 Therapeutic radiology simulation-aided field setting; simple

77285 ; intermediate



77290 ; complex

77295 Three-dimensional radiotherapy plan, including dose-volume histograms

77316 Simple Isodose Planning - Calculation(s) made from one to four sources or remote afterloading brachytherapy, one channel. Includes basic dosimetry calculation(s)

77317 Intermediate Isodose Planning - Calculation(s) made from five to 10 sources, or remote afterloading brachytherapy, two to 12. Includes basic dosimetry calculation(s)

77318 Complex Isodose Planning – Calculation(s) made from more than 10 sources, or remote afterloading brachytherapy, more than 12 channels. Includes basic dosimetry calculation(s)

77331 Special Dosimetry (e.g., TLD, microdosimetry) (specify), only when prescribed by the treating physician

77336 Continuing medical physics consultation, including assessment of treatment parameters, quality assurance of dose delivery and review of patient treatment documentation in support of the radiation oncologist, reported per week of therapy

77370 Special medical radiation physics consultation

77470 Special treatment procedure (e.g., total body irradiation, hemibody radiation, per oral, endocavitary or intraoperative cone irradiation)

The codes used for brachytherapy are categorized in three ways. First, brachytherapy codes are distinguished by the rate at which the dose of radiation is given. There are two major categories:

- Low-dose-rate (LDR) radiation therapy is delivered at a rate less than or equal to 2 Gy/hour at the prescription point.
- High-dose-rate (HDR) radiation therapy is delivered at a dose rate greater than 12 Gy/hour at the prescription point.

The codes can also be categorized by the anatomical position of the applicator relative to the treatment site. There are three categories:

- Interstitial application sources are inserted directly into a tumor, through a surgically created pathway for implant placement, such as breast or prostate.
- Intracavitary (also called intraluminal) application sources are inserted into a body cavity, such as the vagina, uterus, bronchus or esophagus.
- Surface application sources are placed directly on the eye (conjunctiva), skin, breast or other external target surface.

The third dimension to the codes is the procedure's level of complexity. Complexity is determined either by the number of sources, in the case of LDR, or the number of channels, in the case of HDR brachytherapy.

LDR sources can be applied temporarily and removed or can be applied permanently. All HDR brachytherapy is temporary and is sometimes referred to as remote afterloading because it is performed using a computerized robotic delivery device.

77750 Infusion or instillation of radioelement solution (includes 3-month follow-up care)


77761 Intracavitary radiation source application; simple; 1 to 4 sources

77762 ; intermediate; 5 to 10 sources

77763 ; complex; more than 10 intracavitary sources

77778 Interstitial radiation source application, complex, includes supervision, handling, loading of radiation source, when performed; more than 10 interstitial sources

77789 Surface application of low-dose-rate radionuclide source



77790 Supervision, handling, loading of radiation source, do not report in conjunction with 77778

77799 Unlisted procedure, clinical brachytherapy

Q3001 Radioelements for brachytherapy, any type, each

There are brachytherapy-specific codes for HDR radiation treatment delivery. The following levels of complexity are recognized according to the number of channels utilized on the HDR treatment device.

77767 1 channel

Remote afterloading high-dose-rate radionuclide skin surface brachytherapy, includes basic dosimetry, when performed; lesion diameter up to 2.0 cm or 1 channel

77768 2 or more channels

Remote afterloading high-dose-rate radionuclide skin surface brachytherapy, includes basic dosimetry, when performed; lesion diameter over 2.0 cm and 2 or more channels, or multiple lesions

77770 1 channel

Remote afterloading high-dose-rate interstitial or intracavitary radionuclide brachytherapy; includes basic dosimetry, when performed, 1 channel

77771 2-12 channels

Remote afterloading high-dose-rate interstitial or intracavitary radionuclide brachytherapy; includes basic dosimetry, when performed, 2 to 12 channels

77772 over 12 channels

Remote afterloading high-dose-rate interstitial or intracavitary radionuclide brachytherapy; includes basic dosimetry, when performed, more than 12 channels

PROCEDURE GUIDANCE CODES

55860 Exposure of prostate, any approach, for insertion of radioactive substance

76872 Ultrasound, transrectal

76873 Echography, transrectal; prostate volume study for brachytherapy treatment planning (separate procedure)

76965 Ultrasonic guidance for interstitial radioelement application

76000 Fluoroscopy (separate procedure), up to one-hour physician time, other than 71023 or 71034 (eg, cardiac fluoroscopy)

76001 Fluoroscopy, physician time more than one hour, assisting a non-radiologic physician (eg, nephrostolithotomy, ERCP, bronchoscopy, transbronchial biopsy)

76641 Ultrasound, breast, unilateral, real time with image documentation, including axilla when performed; complete

76642 Ultrasound, breast, unilateral, real time with image documentation, including axilla when performed; limited

76942 Ultrasonic guidance for needle placement (e.g., biopsy, aspiration, injection, localization device), imaging supervision and interpretation

76965 Ultrasound guidance for the interstitial radioelement application



77002 Fluoroscopic guidance for needle placement (e.g., biopsy, aspiration, injection, localization device)

77012 Computed tomography guidance for needle placement (e.g., biopsy, aspiration, injection, localization device), radiological supervision and interpretation

77014 Computed tomography guidance for placement of radiation therapy fields

77021 Magnetic resonance guidance for needle placement

Anatomical Region: Skin

17999 Unlisted procedure, skin, mucous membrane and subcutaneous tissue

Anatomical Region: Breast

19296 Placement of radiotherapy afterloading expandable catheter (single or multichannel) into the breast for interstitial radioelement application following partial mastectomy, includes imaging guidance; on date separate from partial mastectomy

19297 Placement of radiotherapy afterloading expandable catheter (single or multichannel) into the breast for interstitial radioelement application following partial mastectomy, includes imaging guidance; concurrent with partial mastectomy (List separately in addition to code for primary procedure) (Use 19297 in conjunction with 19301 or 19302)

19298 Placement of radiotherapy afterloading brachytherapy catheters (multiple tube and button type) into the breast for interstitial radioelement application following (at the time of or subsequent to) partial mastectomy, includes imaging guidance

Anatomical Region: Musculoskeletal

20555 Placement of needles or catheters into muscle and/or soft tissue for subsequent interstitial radioelement application (at the time of or subsequent to the procedure)

Anatomical Region: Trachea and Bronchi

31627 Bronchoscopy, rigid or flexible, including fluoroscopic guidance, when performed; diagnostic, with cell washing, when performed (separate procedure) with computer-assisted, image-guided navigation

Anatomical Region: Lung

31643 Bronchoscopy (rigid or flexible); with placement of catheter(s) for intracavitary radioelement application (For intracavitary radioelement application, see 77761-77763, 77770, 77771, 77772)

32553 Placement of interstitial device(s) for radiation therapy guidance (e.g., fiducial markers, dosimeter), percutaneous, intrathoracic, single or multiple

Anatomical Region: Head and Neck (including Brain)

41019 Placement of needles, catheters, or other device(s) into the head and/or neck region (percutaneous, transoral, or transnasal) for subsequent interstitial radioelement application

0190T Placement of intraocular radiation source applicator

Anatomical Region: Upper Gastrointestinal

43241 Upper gastrointestinal endoscopy including esophagus, stomach and either the duodenum and/or jejunum as appropriate; with transendoscopic intraluminal tube or catheter placement

Anatomical Region: Abdomen, Peritoneum, and Omentum

49411 Placement of interstitial device(s) for radiation therapy guidance (e.g., fiducial markers, dosimeter), percutaneous, intra-abdominal, intra-pelvic (except prostate) and/or retroperitoneum, single or multiple

55920 Placement of needles or catheters into pelvic organs and/or genitalia (except prostate) for subsequent interstitial radioelement application

57155 Insertion of uterine tandem and/or vaginal ovoids for clinical brachytherapy

57156 Insertion of a vaginal radiation afterloading apparatus for clinical brachytherapy

58346 Insertion of Heyman capsules for clinical brachytherapy

58999 Unlisted procedure, female genital system (nonobstetrical)

Anatomical Region: Prostate

55875 Transperineal placement of needles or catheters into prostate for interstitial radioelement application, with or without cystoscopy.

MEDICARE HOSPITAL OUTPATIENT CODES FOR SOURCES AND DEVICES

When hospitals bill Medicare for services performed in the hospital outpatient setting, Medicare will make payment for the LDR source in addition to the treatment delivery.

C1715 Brachytherapy needle

C1717 Brachytherapy source, non-stranded, high-dose-rate iridium-192, per source

C1728 Catheter, brachytherapy seed administration

C1716 Brachytherapy source, non-stranded, gold-198, per source

C1717 Brachytherapy source, non-stranded, high-dose-rate iridium-192, per source

C1719 Brachytherapy source, non-stranded, non-high-dose-rate iridium-192, per source

C2616 Brachytherapy source, non-stranded, yttrium-90, per source

C2634 Brachytherapy source, non-stranded, high activity, iodine-125, greater than 1.01 mCi (NIST), per source

C2635 Brachytherapy source, non-stranded, high activity, palladium-103, greater than 2.2 mCi (NIST), per source

C2636 Brachytherapy linear source, non-stranded, palladium-103, per 1 mm

C2637 Brachytherapy source, non-stranded, ytterbium-169, per source

C2638 Brachytherapy source, stranded, iodine-125, per source

C2639 Brachytherapy source, non-stranded, iodine-125, per source

C2640 Brachytherapy source, stranded, palladium-103, per source

C2641 Brachytherapy source, non-stranded, palladium-103, per source

C2642 Brachytherapy source, stranded, cesium-131, per source

C2643 Brachytherapy source, non-stranded, cesium-131, per source

C2645 Brachytherapy planar source, palladium-103, per square millimeter

C2698 Brachytherapy source, stranded, not otherwise specified, per source

C2699 Brachytherapy source, non-stranded, not otherwise specified, per source

A9527 Iodine I-125, sodium iodide solution, therapeutic, per millicurie

CMS' National Policy:

- Title XVIII of the Social Security Act, section 1862 (a)(7)
 - This section excludes routine physical checkups.
- Title XVIII of the Social Security Act, section 1862 (a)(1)(A)
 - This section states that no Medicare payment shall be made for items or services that are not reasonable and necessary for the diagnosis or treatment of illness or injury.
- Title XVIII of the Social Security Act, section 1833 (e)
 - This section prohibits Medicare payment for any claim that lacks the necessary information for processing.
- Medicare Carrier's Manual, section 15022 (B) (1 and 2)
 - This section prohibits payment of the technical component of radiological procedures by Medicare Part B to hospital inpatients.
- Medicare Carrier's Manual, section 15022 (D)(2 and 4)
 - This section describes the services that are bundled with brachytherapy procedures.



INDICATIONS AND LIMITATIONS OF COVERAGE AND/OR MEDICAL NECESSITY

Indications For Coverage

Brachytherapy may be indicated as a primary or adjunctive therapy in a variety of tumors. A dose rate is selected based on the individual needs of the patient. LDR (low-dose-rate) and HDR (high-dose-rate) brachytherapy are two delivery systems for brachytherapy that use radioactive material to deliver a dose of intensive radiation therapy to a specific well-defined local site (treatment volume). In both LDR and HDR the treatment site should be defined and accessible to the applicators that are the delivery medium for the radioactive sources. This is done to treat a primary or metastatic neoplasm, while sparing sensitive, adjacent normal tissues. LDR and HDR procedures may be given with intent to cure, intent to palliate or to obtain local control (either cure or palliation). Both may be given in conjunction with a course of external beam radiation therapy, or as single modalities.

- LDR is usually delivered over one to several days in a hospital setting; however, LDR may consist of permanently implanted sources and be performed as either an ambulatory or in-patient procedure that delivers radiation as the isotope decays.
- HDR is performed by using a remote afterloading device to deliver the radioactive source(s). HDR allows the dose to be delivered customarily in minutes and usually on an outpatient basis and is often given in a series of multiple fractions.
- Pulsed Dose Rate (PDR) Brachytherapy: Uses sources of intermediate strength and delivers a series of doses on a 1-2 hourly schedule over a 1-2 day treatment period. It is also a form of remote afterloading.

Limitations of Coverage

- Although radiographs may be used in brachytherapy simulation, these images should not be reported as port-films.
- Follow-up visits for 90 days after treatment are not separately payable for CPT codes 77750 and 77761-3. A 90-day global period also effectively applies to HDR treatment (CPT codes 77767-8 and 77770-2) when Radiation Treatment Management (CPT code 77427) is reported during the HDR treatment course. CPT codes 77778 and 77789 are not subject to 90-day global period payment restrictions.
- Only a physician authorized as an authorized user by the Nuclear Regulatory Commission or an Agreement State for brachytherapy should work with radioactive materials.
- Products used for the patient's comfort may not be charged as treatment devices (e.g., pillows, pads or cushions).

ICD-10-CM Codes that may be Associated with Medical Necessity

Note: Diagnosis codes are based on the current ICD-10-CM codes that are effective at the time of the Model Policy publication. Any updates to ICD-10-CM codes will be reviewed by ASTRO, and coverage should not be presumed until the results of such review have been published/posted. These ICD diagnosis codes may support medical necessity under this Model Policy.

Regarding non-melanomatous skin cancers, a distinction should be made between electronic brachytherapy and radionuclide brachytherapy. Radionuclide brachytherapy is a historical treatment modality for skin cancers, and a variety of effective treatment techniques and dose schedules have been developed and used for both LDR and HDR treatments. This form of brachytherapy is particularly useful for patients that may benefit from a shorter treatment course or more precise dosimetry that this modality allows as compared to external beam techniques and should be considered a standard radiotherapy option. Electronic brachytherapy, however, has been more recently applied to skin cancer treatment for which sufficient outcomes data are accruing.

TRUNCATED DIAGNOSIS CODES ARE NOT ACCEPTABLE.

System	Site	ICD-10 Codes
Head and Neck	Lip	C00.0 - C00.9
	Tongue	C01 - C02.9
	Major salivary glands	C07 - C08.9
	Gum	C03.0 - C03.9
	Floor of mouth	C04.0 - C04.9
	Other parts of the mouth	C05.0 - C06.9
	Oropharynx	C09.0 - C10.9
	Nasopharynx	C11.0 - C11.9
	Hypopharynx	C12 - C13.9
	Nasal cavities, middle ear and accessory sinuses	C30.0 - C31.9
	Larynx	C32.0 - C32.9
Digestive Organs and Peritoneum	Esophagus	C15.3 - C15.9
	Stomach	C16.0 - C16.9
	Small intestine	C17.0 - C17.9
	Colon	C18.0 - C18.9
	Rectum, rectosigmoid, anus	C19 - C21.8
	Liver, intrahepatic bile ducts	C22.0 - C22.9
	Gallbladder, extrahepatic bile ducts	C23 - C24.9
	Pancreas	C25.0 - C25.9
	Retroperitoneum, peritoneum	C45.1 C48.0 - C48.8
Respiratory and Intrathoracic Organs	Trachea, bronchus and lung	C33 - C34.92
	Pleura	C38.4 C45.0
	Thymus, heart and mediastinum	C37 - C38.8 C45.2
	Stenosis of coronary artery stent	T82.855A
Bone, connective tissue and skin	Bone	C40.00 - C41.9
	Connective and other soft tissue	C47.0 - C49.9
	Skin	C43.0 - C44.99 D03.0 - D03.9
	Kaposi's sarcoma	C46.0 - C46.9
	Merkel cell carcinoma	C4A.0 - C4A.9 D3A.00 - D3A.8 C7B.00 - C7B.1

Breast	Female breast	C50.011 - C50.019 C50.111 - C50.119 C50.211 - C50.219 C50.311 - C50.319 C50.411 - C50.419 C50.511 - C50.519 C50.611 - C50.619 C50.811 - C50.819 C50.911 - C50.919 D05.00 - D05.92
	Male breast	C50.021 - C50.029 C50.121 - C50.129 C50.221 - C50.229 C50.321 - C50.329 C50.421 - C50.429 C50.521 - C50.529 C50.621 - C50.629 C50.821 - C50.829 C50.921 - C50.929

Genitourinary organs	Cervix	C53.0 - C53.9
	Uterus	C55 C54.0 - C54.9
	Ovary and adnexa	C56.1 - C57.4
	Other female genital organs	C51.0 - C52 C57.7 - C57.9
	Prostate	C61
	Testis	C62.00 - C62.90
	Penis and other male genital organs	C60.0 - C60.9 C63.00 - C63.9
	Bladder	C67.0 - C67.9
Kidney	C64.1 - C66.9 C68.0 - C68.9	

Other sites	Eye	C69.00 - C69.92
	Brain, other parts of nervous system	C70.0 - C72.9
	Endocrine glands	C73 C74.00 - C75.9
	Benign neoplasms of brain, cranial nerves and meninges	D32.0 - D33.3
	Benign neoplasms of pituitary, pineal, aortic body and other paraganglia	D35.2 - D35.6
	Hypertrophic scar	L91.0

Malignant neoplasm of other and ill-defined sites	Various regions	C76.0 - C76.8 C45.7
Secondary and unspecified malignant neoplasm of lymph nodes	Lymph node metastases	C77.0 - C77.9
Secondary malignant neoplasm of respiratory, digestive and other specified sites	Metastatic disease other than lymph node metastases	C78.00 - C80.1 C45.9
Lymphatic and hematopoietic tissue	Non-Hodgkin's lymphoma	C82.00 - C86.6 C91.40 - C91.42 C96.A C96.0 - C96.9 C96.Z
	Hodgkin's lymphoma	C81.00 - C81.99
	Multiple myeloma	C90.00
Neoplasms of uncertain behavior		Need specific sites
Nervous System		C00- D49

Limitations of Coverage

Brachytherapy is not considered reasonable and medically necessary unless at least one of the criteria listed in the "Indications of Coverage" section of this policy is present.

- A claim for brachytherapy submitted without a valid ICD-10-CM diagnosis code will be returned as an incomplete claim under 1833 (e).
- A claim for brachytherapy submitted without one of the ICD-10-CM diagnoses codes listed in the "ICD-10-CM Diagnosis Codes That Support Medical Necessity" section of this policy.
- CPT Evaluation and Management (E/M) codes are available for use by the physician when seeing new patients in the office, the freestanding clinic, hospital setting or ambulatory service center (ASC).
- The physician's professional component for the brachytherapy procedure includes any necessary hospital admission and hospital care during the time that the patient is undergoing the brachytherapy procedure. Admission, subsequent hospital care and discharge day summary is included in the global fee for brachytherapy procedure.
- When E/M services are performed on the same day as brachytherapy, a -25 modifier should accompany the E/M consultation code to reflect that a separate E/M service was provided on the same day.

Documentation Requirements

Documentation in the patient's medical records must support:

- Documentation supporting the medical necessity of these services, such as ICD-10-CM codes, must be submitted with each claim.
- The treatment goal (curative, palliative or tumor control) must be documented in the medical record.
- The record must contain documentation of the patient's informed consent to treatment.
- A written, signed and dated prescription or treatment plan designed by the radiation oncologist must be on file. The prescription must include the following information: designation of the treatment site, designation of the isotope, designation of the number of source positions, and the planned dose to selected points described during dosimetry.
- Given the multiplicity of services that are inherent in brachytherapy, it is essential that the medical records reflect each service in a clear linear and temporally logical form. Flow charts, where helpful, are recommended. All procedures should be documented with a procedural note. A treatment summary should be prepared.
- Since HDR treatments are typically given as a series (often twice daily, over a period of days or weeks) they should be individually documented.


REFERENCES

The medical literature regarding brachytherapy is extensive. The following list comprises a compilation of selected peer reviewed publications reporting clinical outcomes in patients treated with brachytherapy organized by disease site.

Bladder

1. Scher HI, Shipley WU, Herr HW. Cancer of the bladder. In: *Cancer: Principles and Practice of Oncology*. 5th ed. VT DeVita Jr, S Hellman, SA Rosenberg, eds. Philadelphia, PA: Lippincott-Raven; 1997:1315.

Brain

1. Chan TA, Weingart JD, Parisi M, et al. Treatment of recurrent glioblastoma multiforme with GliaSite brachytherapy. *Int J Radiat Oncol Biol Phys*. 2005; 62:1133-1139.
2. Gabayan AJ, Green SB, Sanan A, et al. GliaSite brachytherapy for treatment of recurrent malignant gliomas: A retrospective multi-institutional analysis. *Neurosurgery* 2006;58(4):701-709
3. McDermott MW, Snee PK, Gutin PH. Interstitial brachytherapy for malignant brain tumors. *Semin Surg Oncol*. 1998; 14(1):79-87.
4. Rogers LR, Rock JP, Sillis AK, et al. Results of a phased II trial of the GliaSite radiation therapy system for the treatment of newly diagnosed, resected single brain metastases. *J Neurosurg*. 2006; 105(3):375-384.
5. Sneed PK, McDermott MW, Gutin PH. Interstitial brachytherapy procedures for brain tumors. *Semin Surg Oncol*. 1997; 13(3):157-166.
6. Welsh J, Sanan A, Gabayan AJ, et al. GliaSite brachytherapy boost as part of initial treatment of glioblastoma multiforme: A retrospective multi-institutional pilot study. *Int J Radiat Oncol Biol Phys*. 2007; 68(1):159-165.
7. Wernicke AG, Sherr DL, Schwartz TH, et al. Feasibility and safety of GliaSite brachytherapy in treatment of CNS tumors following neurosurgical resection. *J Can Res Ther*. 2010; 6:65-74.

Breast

1. *Accelerated Partial Breast Irradiation: Techniques and Clinical Implementation*. (Eds) David Wazer MD, Douglas W Arthur MD, Frank Vicini MD, Springer-Verlag 2006
2. Arthur D. Accelerated partial breast irradiation: a change in treatment paradigm for early stage breast cancer. *J Surg Oncol*. 2003; 84(4):185-191.
3. Arthur DW, Koo D, Zwicker RD, et al. Partial breast brachytherapy after lumpectomy: low-dose-rate and high-dose-rate experience. *Int J Radiat Oncol Biol Phys*. 2003; 56(3):681-689.
4. Arthur DW, Vicini FA. Accelerated Partial Breast Irradiation as a part of Breast Conservation Therapy. *J Clin Onc*. 2005; 23(8):1726-1735.
5. Arthur DW, Vicini FA, Kuske RR, et al. Accelerated partial breast irradiation: an updated report from the American Brachytherapy Society. *Brachytherapy*. 2002; 1(4):184-190.
6. *ASTRO/ACR Guide to Radiation Oncology Coding 2010*.
7. *ACR-ASTRO Practice Guideline for the Performance of High-Dose-Rate Brachytherapy Revised 2010*
8. *ACR-ASTRO Practice Guideline for the Performance of Low-Dose-Rate Brachytherapy Revised 2010*
9. Brenner, DJ; Huang, Y; Hall, EJ. Fractionated High Dose-Rate Versus Low Dose-Rate Regimens for Intracavitary Brachytherapy of the Cervix: Equivalent Regimens for Combined Brachytherapy and External Irradiation, *Int J Radiation Oncology Biol Phys*. 1991; 21(6): 1415-1423.
10. Chen PY, Vicini FA, Benitez P, et al. Long-term cosmetic results and toxicity after accelerated partial-breast irradiation: a method of radiation delivery by interstitial brachytherapy for the treatment of early-stage breast carcinoma. *Cancer*. 2006; 106(5):991-999.
11. Cheng, BS; Nagalingam, S; Komanduri, MA; et al. Dosimetric Considerations of Stereotactic Brain Implants, *Int J Radiation Oncology Biol Phys*. 1989; 17(4): 887-891.
12. Consensus statement for Accelerated Partial Breast Irradiation; the American Society of Breast Surgeons.
13. Goyal S, Vicini F, Beitsch PD, et al. Ductal carcinoma in situ treated with breast-conserving surgery and accelerated partial breast irradiation: comparison of the Mammosite registry trial with intergroup study E5194. *Cancer*. 2011; 117(6):1149-1155.

14. Haffty BG, Vicini FA, Beitsch P, et al. Timing of Chemotherapy after MammoSite radiation therapy system breast brachytherapy: Analysis of the American Society of Breast Surgeons MammoSite breast brachytherapy registry trial. *Int J Radiat Oncol Biol Phys.* 2008; 72(5):1441-1448.
15. Harper JL, Watkins JM, Zauls AJ. Six-year experience: Long-term disease control outcomes for partial breast irradiation using MammoSite balloon brachytherapy. *Am J Surg.* 2010; 199(2):204-209.
16. Keisch M, Vicini F, Kuske RR. Two-year outcome with the Mammosite breast brachytherapy applicator: factors associated with optimal cosmetic results when performing partial breast irradiation. *Int J Radiat Oncol Biol Phys.* 2003; 57(2) (Supplement 1):s315.
17. Keisch M, Vicini F, Kuske RR, et al. Initial clinical experience with the MammoSite breast brachytherapy applicator in women with early-stage breast cancer treated with breast-conserving therapy. *Int J Radiat Oncol Biol Phys.* 2003; 55(2):289-293.
18. Khan AJ, Vicini F, Beitsch P, et al. Local control, toxicity, and cosmesis in women younger than 50 enrolled onto the American Society of Breast Surgeons MammoSite Radiation Therapy System registry trial. *Ann Surg Oncol.* 2009; 16(6):1612-8.
19. King TA, Bolton JS, Kuske RR, et al. Long-term results of wide-field brachytherapy as the sole method of radiation therapy after segmental mastectomy for T is,1,2 breast cancer. *Am J Surg.* 2000; 180:299-304.
20. Lawenda BD, Taghian AG, Kachnic LA, et al. Dose-volume analysis of radiotherapy for T1N0 invasive breast cancer treated by local excision and partial breast irradiation by low-dose-rate interstitial implant. *Int J Radiat Oncol Biol Phys.* 2003; 56(3):671-680.
21. McHaffie DR, Patel RR, Adkison JB, et al. Outcomes After Accelerated Partial Breast Irradiation in Patients with ASTRO Consensus Statement Cautionary Features. *Int J Radiat Oncol Biol Phys.* 2011; 81(1): 46-51.
22. Nag S, Kuske RR, Vicini F, et al. The American Brachytherapy Society recommendations for brachytherapy for carcinoma of the breast. *Oncology* 2001; 15(2): 195-205.
23. National Comprehensive Cancer Network® (NCCN). NCCN GUIDELINES™ Clinical Guidelines in Oncology™. © National Comprehensive Cancer Network, Inc 2010, All Rights Reserved. Breast Cancer. V.2.2010. Accessed August 2010. Available at URL address: http://www.nccn.org/professionals/physician_gls/PDF/breast.pdf
24. Nori, D; Allison, R; Kaplan, B; et al. High Dose-Rate Intraluminal Irradiation in Bronchogenic Carcinoma, *Chest.* 1993; 104(4): 1006-1011.
25. Ott OJ, Hildebrandt G, Potter R, et al. Accelerated partial breast irradiation with multi-catheter brachytherapy: Local control, side effects and cosmetic outcome for 274 patients. Results of the German-Austrian multi-centre trial. *Radiation Oncol.* 2007; 82(3):281-286.
26. Polgar C, Major T, Fodor J, et al., High-dose-rate brachytherapy alone versus whole breast radiotherapy with or without tumor bed boost after breast-conserving surgery: seven-year results of a comparative study. *Int J Radiat Oncol Biol Phys.* 2004; 60(4):1173-1181.
27. Polgar C, Strnad V, Major T. Brachytherapy for partial breast irradiation: the European experience. *Semin Radiat Oncol.* 2005; 15(2):116-122.
28. Polgar C, Sulyok Z, Fodor J, et al. Sole brachytherapy of the tumor bed after conservative surgery for T1 breast cancer: five-year results of a phase I-II study and initial findings of a randomized phase III trial. *J Surg Oncol.* 2002; 80(3):121-128; discussion 129.
29. Scanderbeg D, Yashar C, White G, et al. Evaluation of three APBI techniques under NSABP B-39 guidelines. *J Appl Clin Med Phys* 2009; 11(1):3021.
30. Shaitelman SF, Vicini FA, Beitsch P, et al. . Five-year outcome of patients classified using the American Society for Radiation Oncology consensus statement guidelines for the application of accelerated partial breast irradiation: an analysis of patients treated on the American Society of Breast Surgeons MammoSite Registry Trial. *Cancer.* 2010; 116(20):4677-4685.
31. Smith BD, Arthur DW, Buchholz TA, et al. Accelerated partial breast irradiation consensus statement from the American Society for Radiation Oncology (ASTRO). *J Am Coll Surg.* 2009; 209(2):269-277.
32. Strnad V, Hildebrandt G, Potter R, et al. Accelerated partial breast irradiation: 5-year results of the German-Austrian multicenter phase II trial using interstitial multicatheter brachytherapy alone after breast-conserving surgery. *Int J Radiat Oncol Biol Phys.* 2011;80(1):17-24.
33. Strnad V, Ott O, Potter R, et al. Interstitial brachytherapy alone after breast conserving surgery: interim results of a German-Austrian multicenter phase II trial. *Brachytherapy.* 2004; 3(3):115-119.
34. Vicini FA, Antonucci JV, Wallace M, et al. Long-term efficacy and patterns of failure after accelerated partial breast irradiation: a molecular assay-based clonality evaluation. *Int J Radiat Oncol Biol Phys.* 2007; 68(2):341-346.

35. Vicini F, Arthur D, Wazer D, et al. Limitations of the American Society of Therapeutic Radiology and Oncology Consensus Panel guidelines on the use of accelerated partial breast irradiation. *Int J Radiat Oncol Biol Phys.* 2011; 79(4):977-984.
36. Vicini FA, Arthur DW. Breast Brachytherapy: North American Experience. *Seminars of Radiation Oncology.* 2005; 15(2):108-15.
37. Vicini F, Beitsch P, Quiet C, et al. Five-Year Analysis of Treatment Efficacy and Cosmesis by the American Society of Breast Surgeons Mammosite Breast Brachytherapy Registry Trial in Patients Treated with Accelerated Partial Breast Irradiation. *Int J Radiat Oncol Biol Phys.* 2011; 79(3): 808-817.
38. Vicini F, Beitsch PD, Quiet CA, et al. Three-year analysis of treatment efficacy, cosmesis, and toxicity by the American Society of Breast Surgeons MammoSite Breast Brachytherapy Registry Trial in patients treated with accelerated partial breast irradiation (APBI). *Cancer.* 2008; 112(4):758-766.
39. Vicini FA, Kestin L, Chen P, et al. Limited-field radiation therapy in the management of early-stage breast cancer. *J Natl Cancer Inst* 2003; 95(16):1205-1210.
40. Wazer D, Berle L, Graham R, et al. Preliminary results of a phase I/II study of HDR brachytherapy alone for T1/T2 breast cancer. *Int J Radiat Oncol Biol Phys.* 2002; 53(4):889-897.
41. Wazer DE, Lowther D, Boyle T, et al. Clinically evident fat necrosis in women treated with high-dose-rate brachytherapy alone for early-stage breast cancer. *Int J Radiat Oncol Biol Phys.* 2001; 50(1):107-111.
42. Yashar C, Scanderbeg D, Kuske R, et al. Initial Clinical Experience With the Strut-Adjusted Volume Implant (SAVI) Breast Brachytherapy Device for Accelerated Partial-Breast Irradiation (APBI): First 100 Patients With More Than 1 Year of Follow-Up. *Int J Radiat Oncol Biol Phys.* 2011; 80(3): 765-770.

Cervix

1. Demanes DJ, Rodriguez RR, Bendre DD, et al. High-dose-rate Transperineal Interstitial Brachytherapy for Cervical Cancer: High Pelvic Control and Low Complication Rates. *Int J Radiat Oncol Biol Phys.* 1999; 45;11:105-112
2. Erickson B, Eifel P, Moughan J, et al. Patterns of brachytherapy practice for patients with carcinoma of the cervix (1996 – 1999): a patterns of care study. *Int J Radiat Biol Phys.* 2005; 63(4):1083-1092.
3. Lertsanguansinchai P, Lertbutsayanukul C, Shotelersuk K, et al. Phase III randomized trial comparing LDR and HDR brachytherapy in treatment of cervical carcinoma. *Int J Radiat Oncol Biol Phys.* 2004; 59(5):1424-1431.
4. Nag S, Chao C, Erickson B, et al. The American Brachytherapy Society recommendations for low-dose-rate brachytherapy for carcinoma of the cervix. *Int J Radiat Oncol Biol Phys.* 2002; 52(1):33-48.
5. Nag S, Erickson B, Thomadsen B, et al. The American Brachytherapy Society Recommendations for High-Dose-Rate Brachytherapy for Carcinoma of the Cervix. *Int J Radiat Oncol Biol Phys.* 2000; 48(1):201-211
6. Shakespeare TP, Lim KH, Lee KM, et al. American Brachytherapy Society. Phase II study of the American Brachytherapy Society guidelines for the use of high-dose rate brachytherapy in the treatment of cervical carcinoma: Is 45-50.4 Gy radio-chemotherapy plus 31.8 Gy in six fractions high-dose rate brachytherapy tolerable? *Int J Gynecol Cancer* 2006; 16(1):277-282.
7. Toita T, Kato S, Niibe Y, et al. Prospective Multi-Institutional Study of Definitive Radiotherapy With High-Dose-Rate Intracavitary Brachytherapy in Patients With Nonbulky (<4-cm) Stage I and II Uterine Cervical Cancer (JAROG0401/JROSG04-2). *Int J Radiat Oncol Biol Phys.* 2011 Apr 4. [Epub ahead of print]
8. Viani GA, Manta GB, Stefano EJ, et al. Brachytherapy for cervix cancer: Low-dose rate or high-dose rate brachytherapy - a meta-analysis of clinical trials. *J Exp Clin Cancer Res.* 2009; 28:47.
9. Viswanathan AN, Cormack R, Rawal B, et al. Increasing brachytherapy dose predicts survival for interstitial and tandem-based radiation for stage IIIB cervical cancer. *Int J Gynecol Cancer.* 2009; 19(8): 1402-1406.
10. Viswanathan AN, Creutzberg CL, Craighead P, et al. International Brachytherapy Practice Patterns: A Survey of the Gynecologic Cancer Intergroup (GCIg). *Int J Radiat Oncol Biol Phys.* 2010; epub, Dec 22.

Colon and Rectum

1. Cohen AM, Minsky BD, Schilsky RL. Cancer of the rectum. In: *Cancer: Principles and Practice of Oncology.* 5th ed. VT DeVita Jr, S Hellman, SA Rosenberg, eds. Philadelphia, PA: Lippincott-Raven; 1997:1228-1229.
2. Corner C, Bryant L, Chapman C, et al. . High-dose-rate afterloading intraluminal brachytherapy for advanced inoperable rectal carcinoma. *Brachytherapy.* 2010; 9(1):66-70.
3. Grimard L, Stern H, Spaans JN. Brachytherapy and local excision for sphincter preservation in T1 and T2 rectal cancer. *Int J Radiat Oncol Biol Phys.* 2009; 74(3):803-809.

4. Shank B, Cunningham JD, Kelsen, DP. Cancer of the anal region. In: *Cancer: Principles and Practice of Oncology*. 5th ed. VT DeVita Jr, S Hellman, SA Rosenberg, eds. Philadelphia, PA: Lippincott-Raven; 1997:1244-1245.

Esophagus

1. Gaspar LE, Nag S, Herskovic A, et al. American Brachytherapy Society (ABS) consensus guidelines for brachytherapy of esophageal cancer. Clinical Research Committee, American Brachytherapy Society, Philadelphia, PA. *Int J Radiat Oncol Biol Phys*. 1997; 38(1):127-132.
2. Ishikawa H, Nonaka T, Sakurai H, et al. Usefulness of intraluminal brachytherapy combined with external beam radiation therapy for submucosal esophageal cancer: long-term follow-up results. *Int J Radiat Oncol Biol Phys*. 2010; 76(2):452-459.
3. Nemoto K, Yamada S, Nishio M, et al. Results of radiation therapy for superficial esophageal cancer using the standard radiotherapy method recommended by the Japanese Society of Therapeutic Radiology and Oncology (JASTRO) Study Group. *Anticancer Res* 26 (2006), 26(2B) 1507–1512.
4. Sai H, Mitsumori M, Araki N, et al. Long-term results of definitive radiotherapy for stage I esophageal cancer. *Int J Radiat Oncol Biol Phys*. 2005; 62(5): 1339–1344.
5. Tamaki T, Ishikawa H, Takahashi T, et al. Comparison of efficacy and safety of low-dose-rate vs. high-dose-rate intraluminal brachytherapy boost in patients with superficial esophageal cancer. *Brachytherapy*. 2011 Jun 21. [Epub ahead of print]

Endometrium

1. Coon D, Beriwal S, Heron DE, et al. High-dose-rate Rotte “Y” applicator brachytherapy for definitive treatment of medically inoperable endometrial cancer: 10-year results. *Int J Radiat Oncol Biol Phys*. 2008;71(3):779-783.
2. Hasbini, A., Haie-Meder C., Morice P, et al. (2002). Outcome after salvage radiotherapy (brachytherapy +/- external) in patients with a vaginal recurrence from endometrial carcinomas. *Radiother Oncol*. 2002; 65(1): 23-28.
3. Ieso PD, Mullassery V, Shrimali R, et al. Image-guided vulvovaginal interstitial brachytherapy in the treatment of primary and recurrent gynecological malignancies. *Brachytherapy*. 2011 Oct 11. [Epub ahead of print]
4. Jhingran, A., Burke T. W., Eifel PJ, Definitive radiotherapy for patients with isolated vaginal recurrence of endometrial carcinoma after hysterectomy. *Int J Radiat Oncol Biol Phys*. 2003 56(5): 1366-1372.
5. Lin, L. L., P. W. Grigsby, Powell MA, et al. Definitive radiotherapy in the management of isolated vaginal recurrences of endometrial cancer. *Int J Radiat Oncol Biol Phys*. 2005; 63(2): 500-504.
6. Nag S, Erickson B, Parikh S, et al. The American Brachytherapy Society Recommendations for High-Dose-Rate Brachytherapy for Carcinoma of the Endometrium. *Int J Radiat Oncol Biol Phys*. 2000;48(3):779-790
7. Nag S, Orton C, Petereit D, et al. The American Brachytherapy Society recommendations for high-dose-rate brachytherapy for carcinoma of the endometrium. *Int J Radiat Oncol Biol Phys*. 2000;48:779-790.
8. Nag, S., Yacoub S., et al. Interstitial brachytherapy for salvage treatment of vaginal recurrences in previously unirradiated endometrial cancer patients. *Int J Radiat Oncol Biol Phys*. 2002; 54(4): 1153-1159.
9. Nout RA, Smit V, Putter H, et al. Vaginal brachytherapy versus pelvic external beam radiotherapy for patients with endometrial cancer of high-intermediate risk (PORTEC-2): An open-label, non-inferiority, randomised trial. *Lancet*. 2010; 375(9717):816-823.
10. Patel MK, Cote ML, Ali-Fehmi, et al. Trends in the Utilization of Adjuvant Vaginal Cuff Brachytherapy and/or External Beam Radiation Treatment in Stage I and II Endometrial Cancer: A Surveillance, Epidemiology, and End-Results Study. *Int J Radiat Oncol Biol Phys*. 2011 Oct 17. [Epub ahead of print]
11. Small, W Jr., Beriwal, S, Demanes, DJ et al. The American Brachytherapy Society Recommendations for Adjuvant Vaginal Cuff Brachytherapy after Hysterectomy. In Press, *Brachytherapy*, 2011.
12. Small W Jr, Erickson B, Kwakwa F, An American Brachytherapy Society survey regarding the practice patterns of post-operative irradiation for endometrial cancer. *Int J Radiat Oncol Biol Phys*. 2005;63(5):1502-1507.
13. Sorbe BG, Smeds AC. Postoperative vaginal irradiation with high-dose-rate afterloading technique in endometrial carcinoma stage I. *Int J Radiat Oncol Biol Phys*. 1990;18(2):305-314.



Eye

1. Leonard KL, Gagne NL, Mignano JE, et al. A 17-year retrospective study of institutional results for eye plaque brachytherapy of uveal melanoma using (125)I, (103)Pd, and (131)Cs and historical perspective. *Brachytherapy*. 2011; 10(4): 331-339.
2. Melia M, Moy CS, Reynolds SM, et al. Quality of life after iodine 125 brachytherapy vs. enucleation for choroidal melanoma: 5-year results from the Collaborative Ocular Melanoma Study: COMS QOLS Report No. 3. *Arch Ophthalmol*. 2006; 124(2):226-238.
3. Sagoo MS, Shields CL, Mashayekhi A, et al. Plaque radiotherapy for juxtapapillary choroidal melanoma: tumor control in 650 consecutive cases. *Ophthalmology*. 2011;118(2):402-407.
4. Sahel JA, Steeves RA, Albert DM. Intraocular melanoma. In: *Cancer: Principles and Practice of Oncology*. 5th ed. VT DeVita Jr, S Hellman, SA Rosenberg, eds. Philadelphia, PA: Lippincott-Raven; 1997:2004-2005.

Head and Neck

1. Puthawala A, Syed AM, Gamie S, et al. Interstitial low-dose-rate brachytherapy as a salvage treatment for recurrent head-and-neck cancers: long-term results. *Int J Radiat Oncol Biol Phys*. 2001; 51(2): 354-362
2. Bollet M, Lapeyre M, Marchal C, et al. Cervical lymph node relapses of head-and-neck squamous cell carcinoma: is brachytherapy a therapeutic option? *Int J Radiat Oncol Biol Phys*. 2001; 51(5), 1305-1312.
3. Gibbs IC, Le QT, Shah RD, et al. Long-term outcomes after external beam irradiation and brachytherapy boost for base-of-tongue cancers. *Int J Radiat Oncol Biol Phys*. 2003; 57(2), 489-494.
4. Han P, Hu K, Culliney B, et al. Concurrent chemoradiation followed by interstitial brachytherapy boost and neck dissection for T4 base of tongue cancer. *Int J Radiat Onco Biol Phys*. 2005; 63(s1):S358
5. Harrison LB, Weissberg JB. A technique for interstitial nasopharyngeal brachytherapy. *Int J Radiat Onco Biol Phys*. 1987; 13(3):451-453
6. Harrison LB, Zelefsky MJ, Pfister DG, et al. Detailed quality of life assessment in patients treated with primary radiotherapy for squamous cell cancer of the base of the tongue. *Head neck*. 1997 19:3 (169-175)
7. Hepel JT, Syed AMN, Puthawala A, et al. Salvage high-dose-rate (HDR) brachytherapy for recurrent head-and-neck cancer. *Int J Radiat Onco Biol Phys*. 2005; 62(5), 1444-1450.
8. Kakimoto N, Inoue T, Inoue T, et al. Results of low- and high-dose-rate interstitial brachytherapy for T3 mobile tongue cancer. *Radiotherapy & Oncology*. 2003; 68(2), 123-128.
9. Lapeyre M, Bollet MA, Racadot S, et al. Postoperative brachytherapy alone and combined postoperative radiotherapy and brachytherapy boost for squamous cell carcinoma of the oral cavity, with positive or close margins. *Head neck*. 2004; 26(3), 216-223.
10. Leung TW, Tung SY, Sze WK, et al. Salvage radiation therapy for locally recurrent nasopharyngeal carcinoma. *Int J Radiat Onco Biol Phys*. 2000; 48(5), 1331-1338.
11. Marsili H, Haie-Mède C, Sassa G, et al. Brachytherapy for T1-T2 floor-of-the-mouth cancers: the Gustave-Roussy Institute experience. *Int J Radiat Onco Biol Phys*. 2002; 52(5), 1257-1263.
12. Nag S, Cano ER, Demanes DJ, et al. The American Brachytherapy Society recommendations for high-dose-rate brachytherapy for head-and-neck carcinoma. *Int J Radiat Oncol Biol Phys*. 2001; 50(5):1190-1198.
13. Pellizzon ACA, Dos Santos Novaes PER, Conte Maia MA, et al. Interstitial high-dose-rate brachytherapy combined with cervical dissection on head and neck cancer. *Head neck*. 2005; 27(12), 1035-1041.
14. Sabbas AM, Kulidzhanov FG, Presser J, et al. HDR brachytherapy with surface applicators: technical considerations and dosimetry. *Technology in cancer research treatment*. 2004; 3(3), 259-267.
15. Spenser S, Harris J, Wheeler R, et al. Late effects of RTOG-9610: Re-irradiation and chemotherapy in patients with squamous cell cancer of the head and neck. *Int J Radiat Onco Biol Phys*. 2003; 57(2 Suppl):S307.
16. Strnad V, Geiger M, Lotter M, et al. The role of pulsed-dose-rate brachytherapy in previously irradiated head-and-neck cancer. *Brachytherapy*. 2002;1(3): 149-153
17. Strnad V, Melzner W, Geiger M, et al. Role of interstitial PDR brachytherapy in the treatment of oral and oropharyngeal cancer. A single-institute experience of 236 patients. *Strahlenther Onkol*. 2005; 181(12): 762-767.
18. Syed AM, Puthawala AA, Damore SJ, et al. Brachytherapy for primary and recurrent nasopharyngeal carcinoma: 20 years experience at Long Beach Memorial. *Int J Radiat Onco Biol Phys*. 2000; 47(5), 1311-1321.



Intracoronary

1. Costa MA, Sabat M, van der Giessen WJ, et al. Late coronary occlusion after intracoronary brachytherapy. *Circulation*. 1999; 100(8):789-792.
2. Waksman R. Intracoronary brachytherapy in the Cath Lab. Physics dosimetry, technology and safety considerations. *Herz*. 1998; 23(6):401-406.

Lung

1. Fernando HC, Landreneau RJ, Mandrekar SJ, et al. The impact of adjuvant brachytherapy with sublobar resection on pulmonary function and dyspnea in high-risk operable patients: preliminary results from the ACOSOG Z4032 trial. *J Thorac Cardiovasc Surg*. 2011; 142(3): 554-562.
2. Fernando HC, Santos RS, Benfield JR, et al. Lobar and sublobar resection with and without brachytherapy for small stage IA non-small cell lung cancer. *J Thorac Cardiovasc Surg*. 2005; 129(2):261-267.
3. Lee W, Daly BD, DiPetrillo TA, et al. Limited resection for non-small cell lung cancer: observed local control with implantation of I-125 brachytherapy seeds. *Ann Thorac Surg*. 2003; 75(1):237-243.
4. Minna JD, Sekido, Y, Fong, K, et al. Cancer of the lung. In: *Cancer: Principles and Practice of Oncology*. 5th ed. VT DeVita Jr, S Hellman, SA Rosenberg, eds. Philadelphia, PA: Lippincott-Raven; 1997:882.
5. Raben A, Mychalczak B. Brachytherapy for non-small cell lung cancer and selected neoplasms of the chest. *Chest*. 1997;112(4 Suppl):276S-286S.
6. Santos R, Colonias A, Parda D, et al. Comparison between sublobar resection and I-125 brachytherapy after sublobar resection in high-risk patients with stage I non-small cell lung cancer. *Surgery*. 2003; 134(4):691-697.
7. Schuchert MJ, Abbas G, Pennathur A, et al. Sublobar resection for early-stage lung cancer. *Semin Thorac Cardiovasc Surg*. 2010 Spring; 22(1):22-31.
8. Tselis N, Ferentinos K, Kolotas C, Schirren J, et al. Computed tomography-guided interstitial high-dose-rate brachytherapy in the local treatment of primary and secondary intrathoracic malignancies. *J Thorac Oncol*. 2011; 6(3):545-552.
9. Voynov G, Heron DE, Lin CJ, et al. Intraoperative (125)I Vicryl mesh brachytherapy after sublobar resection for high-risk stage I non-small cell lung cancer. *Brachytherapy*. 2005; 4(4):278-285.

Keloids

1. Van Leeuwen MC, et al. "High-dose-rate brachytherapy for the treatment of recalcitrant keloids: a unique, effective treatment protocol." *Plast Reconstr Surg*. 2014 Sep;134(3):527-34. doi: 10.1097/PRS.0000000000000415.

Prostate

1. Arvold ND, Chen MH, Moul JW, et al. Risk of death from prostate cancer after radical prostatectomy or brachytherapy in men with low or intermediate risk disease. *J Urol*. 2011; 186(1):91-96.
2. Corner C, Rojas AM, Bryant L, et al. A Phase II study of high-dose-rate afterloading brachytherapy as monotherapy for the treatment of localized prostate cancer. *Int J Radiat Oncol Biol Phys*. 2008; 72(2):441-446.
3. Crook J, Borg J, Evans A, et al. 10-year experience with I-125 prostate brachytherapy at the princess Margaret hospital: results for 1,100 patients. *Int J Radiat Oncol Biol Phys*. 2011; 80:1323-1329.
4. Demanes DJ, Martinez AA, Ghilezan M, et al. High-Dose-Rate Monotherapy: Safe and Effective Brachytherapy for Patients with Localized Prostate Cancer. *Int J Radiat Oncol Biol Phys*. 2011 Feb 10. [Epub ahead of print]
5. Deutsch I, Zelefsky MJ, Zhang Z, et al. Comparison of PSA relapse-free survival in patients treated with ultra-high-dose IMRT versus combination HDR brachytherapy and IMRT. *Brachytherapy* 2010; 9(4):313-318.
6. Keyes M, Spadinger I, Liu M, et al. Rectal toxicity and rectal dosimetry in low-dose-rate iodine-125 permanent prostate implants: A long-term study in 1006 patients. *Brachytherapy*. 2010; 9(Suppl 1): S79-S80.
7. Nag S, Beyer D, Friedland J, et al. American Brachytherapy Society (ABS) recommendations for transperineal permanent brachytherapy of prostate cancer. *Int J Radiat Oncol Biol Phys*. 1999; 44(4):789-799.
8. Rosenthal SA, Bittner NH, Beyer DC, et al. American Society for Radiation Oncology (ASTRO) and American College of Radiology (ACR) practice guideline for the transperineal permanent brachytherapy of prostate cancer. *Int J Radiat Oncol Biol Phys*. 2011; 79(2):335-341.


9. Stock RG, Stone NN. Current topics in the treatment of prostate cancer with low-dose-rate brachytherapy. *Urol Clin North Am.* 2010; 37(1):83-96.
10. Sylvester JE, Grimm PD, Blasko JC, et al. 15-Year biochemical relapse free survival in clinical Stage T1-T3 prostate cancer following combined external beam radiotherapy and brachytherapy; Seattle experience. *Int J Radiat Oncol Biol Phys.* 2007; 67(1):57-64.
11. Taira AV, Merrick GS, Butler WM, et al. Long-term outcome for clinically localized prostate cancer treated with permanent interstitial brachytherapy. *Int J Radiat Oncol Biol Phys.* 2011; 79(5):1336-1342.
12. Thompson A, Keyes M, Pickles T, et al. Evaluating the Phoenix definition of biochemical failure after (125)I prostate brachytherapy: Can PSA kinetics distinguish PSA failures from PSA bounces? *Int J Radiat Oncol Biol Phys.* 2010; 78(2):415-421.
13. Vicini FA, Kini VR, Edmundson G, et al. A comprehensive review of prostate cancer brachytherapy: Defining an optimal technique. *Int J Radiat Oncol Biol Phys.* 1999; 44(3):483-491.
14. Zelefsky MJ. PSA bounce versus biochemical failure following prostate brachytherapy. *Nat Clin Pract Urol.* 2006;3(11):578-579.
15. Zelefsky MJ, Chan H, Hunt M, et al. Long-term outcome of high dose intensity modulated radiation therapy for patients with clinically localized prostate cancer. *J Urol.* 2006; 176(4 pt 1): 1415-1419.
16. Zelefsky MJ, Yamada Y, Cohen GN, et al. Five-year outcome of intraoperative conformal permanent I-125 interstitial implantation for patients with clinically localized prostate cancer. *Int J Radiat Oncol Biol Phys.* 2007; 67(1): 65-70.
17. Zelefsky MJ, Yamada Y, Pei X, et al. Comparison of tumor control and toxicity outcomes of high-dose intensity-modulated radiotherapy and brachytherapy for patients with favorable risk prostate cancer. *Urology.* 2011; 77(4):986-990.

Sarcomas

1. Brennan MF, Casper ES, Harrison LB. Soft tissue sarcoma. In: *Cancer: Principles and Practice of Oncology.* 5th ed. VT DeVita Jr, S Hellman, SA Rosenberg, eds. Philadelphia, PA: Lippincott-Raven; 1997:1762-1764.
2. Hilaris BS, Bodner WR, Mastoras CA. Role of brachytherapy in adult soft tissue sarcomas. *Semin Surg Oncol.* 1997; 13(3):196-203.
3. Nag S, Shasha D, Janjan N, et al. The American Brachytherapy Society recommendations for brachytherapy of soft tissue sarcomas. *Int J Radiat Oncol Biol Phys.* 2001; 49(4):1033-1043.
4. O'Sullivan B, Davis AM, Turcotte R, et al. Preoperative versus postoperative radiotherapy in soft-tissue sarcoma of the limbs: a randomised trial. *Lancet.* 2002; 359 (9325): 2235-2241
5. Pisters PW, Harrison LB, Leung DH, et al. Long-term results of a prospective randomized trial of adjuvant brachytherapy in soft tissue sarcoma. *J Clin Oncol.* 1996; 14(3):859-868.

Skin

1. Alam M, Nanda S, Mittal BB, et al. The use of brachytherapy in the treatment of nonmelanoma skin cancer: A review. *J Am Acad Dermatol.* 2011; 65(2):377-88.
2. Somanchi B, Stanton A, Webb M, et al. Hand function after high-dose-rate brachytherapy for squamous cell carcinoma of the skin of the hand. *Clin Oncol (R Coll Radiol)* 2008; 20(9):691-697.
3. Bhatnagar A, Loper A. The Initial Experience of electronic brachytherapy for the treatment of non-melanoma skin cancer. *Radiation Oncology.* 2010; 5:87.
4. Conill C, Verger E, Marruecos J, et al. Low-dose-rate brachytherapy in lip carcinoma, *Clin Transl Oncol* 2004; 9(4):251-254.
5. Kohler-Brock A, Pragger W, Pohlmann S, et al. The Indications for and results of HDR afterloading therapy in diseases of the skin and mucosa with standardized surface applicators (The Leipzig Applicator), *Strahlenther Onkol* 1999; 175(4): 170-174.
6. Debois J. Cesium-137 brachytherapy for epithelioma of the skin of the nose: experience with 370 patients. *J Belge Radiol.* 1994; 77(1):1-4,
7. Guix B, Finestres F, Tello J, et al. Treatment of Skin Carcinomas of the Face by High-dose-rate Brachytherapy and Custom Made Surface Molds. *Int J Radiat Oncol Biol Phys.* 2000; 47(1): 95-102.
8. Lee JD, Park KK, Lee M, et al. Radionuclide therapy of skin cancers and Bowen's disease using a specially designed skin patch. *J Nucl Med.* 1997, 38(5):697-702.
9. Rio E, Bardet E, Ferron C, et al. Interstitial brachytherapy of periorificial skin carcinomas of the face: a retrospective study of 97 cases, *Int J Radiat Oncol Biol Phys.* 2005; 63(3):753-757.

- 
10. Rudoltz M, Perkins R, Luthmann R, et al. High-dose-rate brachytherapy with a custom-surface mold to treat recurrent squamous cell carcinomas of the skin of the forearm. *J Am Acad Dermatol*. 1998; 38(6 Pt 1):1003-1005.
 11. Sedda A, Rossi G, Cipriani C, et al. Dermatological high-dose-rate brachytherapy for the treatment of basal and squamous cell carcinoma. *Clin Exp Dermatol*. 2008; 33(6):745-749.
 12. Svoboda V, Kovarik J, Morris F. High dose-rate microselectron molds in the treatment of skin tumors. *Int J Radiat Oncol Biol Phys*. 1995; 31(4):967-972.

Vagina

1. Beriwal S, Heron DE, Mogus R, et al. High-dose rate brachytherapy (HDRB) for primary or recurrent cancer in the vagina. *Radiat Oncol*. 2008 Feb 13;3:7.
2. Beriwal S, Rwigema JC, Higgins E, et al. 3D image-based HDR interstitial brachytherapy for vaginal cancer. *Brachytherapy*. 2011 Jun 10. [Epub ahead of print]
3. Dimopoulos JC, Schmid MP, Fidarova E, et al. Treatment of Locally Advanced Vaginal Cancer with Radiochemotherapy and Magnetic Resonance Image-guided Adaptive Brachytherapy: Dose-volume Parameters and First Clinical Results. *Int J Radiat Oncol Biol Phys*. 2011 Aug 24. [Epub ahead of print].
4. Frank SJ, Jhingran A, Levenback C, et al. Definitive radiation therapy for squamous cell carcinoma of the vagina. *Int J Radiat Oncol Biol Phys*. 2005;62(1):138-147.
5. Ieso PD, Mullassery V, Shrimali R, et al. Image-guided vulvovaginal interstitial brachytherapy in the treatment of primary and recurrent gynecological malignancies. *Brachytherapy*. 2011 Oct 11. [Epub ahead of print]
6. Jhingran A, Burke TW, Eifel PJ. Definitive radiotherapy for patients with isolated vaginal recurrence of endometrial carcinoma after hysterectomy. *Int J Radiat Oncol Biol Phys*. 2003;56(5):1366-1372.
7. Kushner DM, Fleming PA, Kennedy AW, et al. High-dose-rate (192)Ir afterloading brachytherapy for cancer of the vagina. *Br J Radiol*. 2003;76(910):719-725.
8. Lieskovsky YE, Demanes D. Combination high-dose Rate brachytherapy and external beam radiation therapy for the treatment of primary vaginal cancer: 5-year results. *Int J Radiat Oncol Biol Phys*. 2004;60(1):S308 [Abstract].
9. Nag S, Yacoub S, Copeland LJ, et al. Interstitial brachytherapy for salvage treatment of vaginal recurrences in previously unirradiated endometrial cancer patients. *Int J Radiat Oncol Biol Phys*. 2002;54(4):1153-1159.
10. Pai HH, Souhami L, Clark BG, et al. Isolated vaginal recurrences in endometrial carcinoma: treatment results using high-dose-rate intracavitary brachytherapy and external beam radiotherapy. *Gynecol Oncol*. 1997;66(2):300-307.
11. Patel MK, Cote ML, Ali-Fehmi, et al. Trends in the Utilization of Adjuvant Vaginal Cuff Brachytherapy and/or External Beam Radiation Treatment in Stage I and II Endometrial Cancer: A Surveillance, Epidemiology, and End-Results Study. *Int J Radiat Oncol Biol Phys*. 2011 Oct 17. [Epub ahead of print]
12. Petignat P, Jolicoeur M, Alobaid A, et al. Salvage treatment with high-dose-rate brachytherapy for isolated vaginal endometrial cancer recurrence. *Gynecol Oncol*. 2006;101(3):445-449.