

Short Term Androgen Deprivation Therapy Without or With Pelvic Lymph Node Treatment Added to Prostate Bed Only Salvage Radiation Therapy: The NRG Oncology/RTOG 0534 SPPORT Trial

A. Pollack¹, T. G. Garrison², A. G. Balogh³, D. Low⁴, D. W. Bruner⁵, J. S. Wefel⁶, L. G. Gomella⁷, E. Vigneault⁸, J. M. Michalski⁹, S. Angyalfi¹⁰, H. Lukka¹¹, S. L. Faria¹², G. Rodrigues¹³, M. C. Beauchemin¹⁴, S. A. Seaward¹⁵, A. M. Allen¹⁶, D. C. Monitto¹⁷, W. Seiferheld², and H. M. Sandler¹⁸

¹University of Miami Sylvester Comprehensive Cancer Center, Miami, FL, ²NRG Oncology SDMC, Philadelphia, PA, ³University of Calgary, Calgary, AB, Canada, ⁴UCLA, Los Angeles, CA, ⁵Nell Hodgson Woodruff School of Nursing, and Winship Cancer Institute at Emory University, Atlanta, GA, ⁶University of Texas MD Anderson Cancer Center, Houston, TX, ⁷Sidney Kimmel Cancer Center of Thomas Jefferson University, Philadelphia, PA, ⁸CHU de Québec, University of Laval, Quebec, QC, Canada, ⁹Washington University School of Medicine, St. Louis, MO, ¹⁰Tom Baker Cancer Centre, Calgary, AB, Canada, ¹¹McMaster University, Hamilton, ON, Canada, ¹²McGill University Health Centre, Montreal, QC, Canada, ¹³London Health Sciences Center, London, ON, Canada, ¹⁴Centre Hospitalier de l'Université de Montréal, Montreal, QC, Canada, ¹⁵Kaiser Permanente, Vallejo, CA, ¹⁶Davidoff Center, Rabin Medical Center, Tel Aviv, Israel, ¹⁷Spartanburg Regional Medical Center, Spartanburg, SC, ¹⁸Cedars Sinai Medical Center, Los Angeles, CA



Disclosure for Dr. Pollack

- Employment
 - University of Miami Miller School of Medicine: Chair and Professor of Radiation Oncology, Deputy Director Sylvester Comprehensive Cancer Center: Employee
- Compensation
 - City of Hope: Honoraria, Travel Expenses; Radiation Therapy Oncology Group: Research Grants, Travel Expenses; Varian: Research Grants; Varian Medical Systems: Research Grants



Background

- Biochemical failure after salvage prostate bed radiation therapy (**PBRT**) is typically 30-40% at 5-10 years.
- Neoadjuvant and concurrent short term androgen deprivation therapy (**STAD**) is an effective addition to primary radiation therapy and in 2005 had not yet been tested with salvage PBRT.
- Pelvic lymph node radiation therapy (**PLNRT**) has shown promise, but has never been conclusively proven to be effective in a Phase III randomized trial.
- Trial hypothesis: For prostate cancer patients with a rising PSA after prostatectomy, there will be an incremental gain in freedom from progression with the addition of:
 - STAD to PBRT
 - PLNRT + STAD to PBRT

NRG Oncology/RTOG 0534/SPPORT Trial Design

S	SV Involvement 1. No 2. Yes	R	Arm 1: PBRT Alone PBRT 64.8-70.2 Gy
T	Prostatectomy Gleason Score 1. Gleason \leq 7 2. Gleason 8-9	A	Arm 2: PBRT + STAD PBRT 64.8-70.2 Gy + STAD for 4-6 months beginning 2 months before RT
R		N	
A		D	
T	Pre-Radiotherapy PSA 1. PSA \geq 0.1 and \leq 1.0 ng/mL 2. PSA > 1.0 and < 2.0 ng/mL	O	Arm 3: PLNRT + PBRT + STAD PLNRT to 45 Gy and PBRT to 64.8-70.2 Gy,+ STAD for 4-6 months beginning 2 months before RT
I		M	
F		I	
Y	Pathology Stage 1. pT2 and margin negative 2. All others	Z	
		E	

SV = seminal vesicle; RT = radiation therapy; PBRT = prostate bed RT; PLNRT = pelvic lymph node RT; STAD = neoadjuvant and concurrent short-term androgen deprivation

Primary endpoint:
FFP at 5 years

Failure defined as first occurrence of:

- $\text{PSA} \geq \text{Nadir}+2 \text{ ng/mL}$
- Clinical progression (local, regional or distant)
- Death due to any cause



Patient characteristics


Patient/Tumor Characteristic	PBRT Alone (n=574)		PBRT+STAD (n=585)		PLNRT+PBRT+STAD (n=577)		Total	
	n	%	n	%	n	%	n	%
Zubrod Performance Status								
0	533	92.9	546	93.3	543	94.1	1622	93.4
1	41	7.1	39	6.7	34	5.9	114	6.6
Seminal Vesicle Involvement								
No	492	85.7	498	85.1	489	84.7	1479	85.2
Yes	82	14.3	87	14.9	88	15.3	257	14.8
Prostatectomy Stage								
pT2	304	53.0	326	55.7	312	54.1	942	54.3
pT3 NOS	19	3.3	17	2.9	22	3.8	58	3.3
pT3a ECE	173	30.1	160	27.4	156	27.0	489	28.2
pT3b SVI	78	13.6	82	14.0	87	15.1	247	14.2
Prostatectomy Margins								
Positive	292	50.9	292	49.9	286	49.6	870	50.1
Negative	273	47.6	286	48.9	288	49.9	847	48.8
Unknown	9	1.6	7	1.2	3	0.5	19	1.1

Median Age=64y; Caucasian 83%; Non-Hispanic 91%



FFP: Interim analysis population (1,191 patients)

Minimum potential FU 5 yr;
Median FU 6.4 yr



No. at Risk	Years from Randomization								
	0	1	2	3	4	5	6	7	8
PBRT Alone	399	363	324	286	259	223	150	84	34
PBRT+NC-STAD	398	381	360	329	299	265	170	91	37
PLNRT+PBRT+NC-STAD	394	385	371	352	334	311	208	125	51

5 yr Rate Comparison

Arm 3 vs Arm 1: $p<0.0001$


Arm 2 vs Arm 1: $p=0.0001$

Arm 3 vs Arm 2: $p=0.0063$

DMC Recommendation

Report findings for all three arms

FFP: All eligible patients (1,792)



No. at Risk	Years from Randomization								
PBRT Alone	573	529	480	417	334	243	165	89	37
PBRT+NC-STAD	585	559	532	467	366	275	179	94	39
PLNRT+PBRT+NC-STAD	576	563	540	488	399	315	209	126	52

5 yr Rate Comparison

Arm 3 vs Arm 1: $p<0.0001$

Arm 2 vs Arm 1: $p<0.0001$

Arm 3 vs Arm 2: $p=0.0039$


HRs and 97.5% CIs

3 vs 1: 0.45 (0.34-0.61)

2 vs 1: 0.62 (0.47-0.82)

3 vs 2: 0.71 (0.52-0.98)

Freedom from distant metastasis: All eligible patients



No. at Risk	Years from Randomization								
PBRT Alone	573	546	522	482	398	302	214	123	54
PBRT+NC-STAD	585	562	547	499	405	315	213	111	48
PLNRT+PBRT+NC-STAD	576	565	551	502	417	330	220	134	56

5 yr Rate Comparison
Arm 3 vs Arm 1: $p=0.014$
Arm 2 vs Arm 1: $p=0.05$
Arm 3 vs Arm 2: $p=0.28$

HRs and 97.5% CIs
3 vs 1: 0.52 (0.30-0.92)
2 vs 1: 0.81 (0.49-1.33)
3 vs 2: 0.64 (0.36-1.14)

No statistically significant differences in OS

Acute Adverse Events (CTCAEv3) Without Regard For Attribution

Type	PBRT Alone (n=557)		PBRT+STAD (n=568)		PLNRT+PBRT+STAD (n=567)		p-value
	n	%	n	%	n	%	
GI							
Grade 2+	11	2.0	22	3.9	39	6.9	<0.001
Grade 3+	1	0.2	5	0.9	4	0.7	0.37
Renal/GU							
Grade 2+	54	9.7	68	12.0	69	12.3	0.35
Grade 3+	5	0.9	5	0.9	8	1.5	0.70
Blood/Bone Marrow							
Grade 2+	13	2.3	10	1.8	29	5.1	0.002
Grade 3+	3	0.5	1	0.2	15	2.6	<0.001



Late Adverse Events (CTCAEv3) Without Regard For Attribution

Type	PBRT Alone (n=557)		PBRT+STAD (n=571)		PLNRT+PBRT+STAD (n=570)		p-value
	n	%	n	%	n	%	
GI							
Grade 2+	54	9.7	50	8.8	46	8.1	0.63
Grade 3+	4	0.7	2	0.4	6	1.1	0.34
Renal/GU							
Grade 2+	188	33.8	184	32.2	197	34.6	0.70
Grade 3+	24	4.4	28	4.9	34	6.0	0.44
Blood/Bone Marrow							
Grade 2+	17	3.1	9	1.6	23	4.1	0.044
Grade 3+	2	0.4	1	0.2	6	1.1	0.12



Conclusions and treatment implications

- Strongest level I evidence supporting PLNRT
- For Arm 3 vs Arm 1, number needed to treat to prevent one progression within 5 years is **6** (95%CI 4.6-8.6)!
 - Follow-up continuing to further elucidate the magnitude of the differences between Arms 2 and 3
- Robust effect translating into a decrease in distant metastasis
- Is there a PSA cutpoint below which PLNRT is not needed?
- Role of PET in PLNRT decisions?
- Implications for the management of primarily managed prostate cancer
 - Local control needed to realize the impact of PLNRT (NRG 0924)

