

Measure Title	Measure Number		Measure Description	High Priority	Data Submission Method(s)			
	NQF	Quality Number (Q#)			Claims	EHR	Registry	QOPI QCDR (by ASCO and ASTRO)
Diabetes: Hemoglobin A1c (HbA1c) Poor Control (>9%)	0059	001	Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period	X	X	X	X	
Medication Reconciliation Post-Discharge	0097	046	The percentage of discharges from any inpatient facility (e.g. hospital, skilled nursing facility, or rehabilitation facility) for patients 18 years and older of age seen within 30 days following discharge in the office by the physician, prescribing practitioner, registered nurse, or clinical pharmacist providing on-going care for whom the discharge medication list was reconciled with the current medication list in the outpatient medical record. This measure is reported as three rates stratified by age group: <ul style="list-style-type: none"> • Submission Criteria 1: 18-64 years of age • Submission Criteria 2: 65 years and older • Total Rate: All patients 18 years of age and older 	X	X	-	X	
Care Plan	0326	047	Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan	X	X	-	X	X
Prostate Cancer: Avoidance of Overuse of Bone Scan for Staging Low Risk Prostate Cancer Patients	0389	102	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at low (or very low) risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy, OR cryotherapy who did not have a bone scan performed at any time since diagnosis of prostate cancer	-	-	X	X	X
Prostate Cancer: Combination Androgen Deprivation Therapy for High Risk or Very High Risk Prostate Cancer	0390	104	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at high or very high risk of recurrence receiving external beam radiotherapy to the prostate who were prescribed androgen deprivation therapy in combination with external beam radiotherapy to the prostate	-	-	-	X	X
Preventive Care and Screening: Influenza Immunization	0041	110	Percentage of patients aged 6 months and older seen for a visit between October 1 and March 31 who received an influenza immunization OR who reported previous receipt of an influenza immunization	-	X	X	X	
Breast Cancer Screening	2372	112	Percentage of women 50-74 years of age who had a mammogram to screen for breast cancer	-	X	X	X	
Colorectal Cancer Screening	0034	113	Percentage of adults 50-75 years of age who had appropriate screening for colorectal cancer	-	X	X	X	
Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan	0421	128	Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous twelve months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter <u>Normal Parameters:</u> Age 18 years and older BMI ≥ 18.5 and < 25 kg/m2	-	X	X	X	
Documentation of Current Medications in the Medical Record	0419	130	Percentage of visits for patients aged 18 years and older for which the eligible professional or eligible clinician attests to documenting a list of current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration	X	X	X	X	X
Pain Assessment and Follow-Up	0420	131	Percentage of visits for patients aged 18 years and older with documentation of a pain assessment using a standardized tool(s) on each visit AND documentation of a follow-up plan when pain is present	X	X	-	X	

Measure Title	NQF	Quality Number (Q#)	Measure Description	High Priority	Measure Category			QOPI QCDR (by ASCO and ASTRO)
					Claims	EHR	Registry	
Preventive Care and Screening: Screening for Depression and Follow-Up Plan	0418	134	Percentage of patients aged 12 years and older screened for depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen	-	X	X	X	X
Melanoma: Continuity of Care – Recall System	0650	137	Percentage of patients, regardless of age, with a current diagnosis of melanoma or a history of melanoma whose information was entered, at least once within a 12 month period, into a recall system that includes: • A target date for the next complete physical skin exam, AND • A process to follow up with patients who either did not make an appointment within the specified timeframe or who missed a scheduled appointment	X	-	-	X	
Melanoma: Coordination of Care	N/A	138	Percentage of patient visits, regardless of age, with a new occurrence of melanoma that have a treatment plan documented in the chart that was communicated to the physician(s) providing continuing care within one month of diagnosis	X	-	-	X	
Oncology: Medical and Radiation – Pain Intensity Quantified	0384	143	Percentage of patient visits, regardless of patient age, with a diagnosis of cancer currently receiving chemotherapy or radiation therapy in which pain intensity is quantified	X	-	X	X	X
Oncology: Medical and Radiation – Plan of Care for Pain	0383	144	Percentage of visits for patients, regardless of age, with a diagnosis of cancer currently receiving chemotherapy or radiation therapy who report having pain with a documented plan of care to address pain	X	-	-	X	
Falls: Risk Assessment	0101	154	Percentage of patients aged 65 years and older with a history of falls that had a risk assessment for falls completed within 12 months	X	X	-	X	
Falls: Plan of Care	0101	155	Percentage of patients aged 65 years and older with a history of falls that had a plan of care for falls documented within 12 months	X	X	-	X	
Oncology: Radiation Dose Limits to Normal Tissues	0382	156	Percentage of patients, regardless of age, with a diagnosis of breast, rectal, pancreatic or lung cancer receiving 3D conformal radiation therapy who had documentation in medical record that radiation dose limits to normal tissues were established prior to the initiation of a course of 3D conformal radiation for a minimum of two tissues	-	X	-	X	
Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention	0028	226	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received tobacco cessation intervention if identified as a tobacco user	-	X	X	X	X
Controlling High Blood Pressure	0018	236	Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90mmHg) during the measurement period	X	X	X	X	
Use of High-Risk Medications in the Elderly	0022	238	Percentage of patients 65 years of age and older who were ordered high-risk medications. Two rates are submitted. 1) Percentage of patients who were ordered at least one high-risk medication. 2) Percentage of patients who were ordered at least two of the same high-risk medication	X	-	X	X	
Cervical Cancer Screening	0032	309	Percentage of women 21-64 years of age who were screened for cervical cancer using either of the following criteria: * Women age 21-64 who had cervical cytology performed every 3 years * Women age 30-64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years	-	-	X	-	
Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented	N/A	317	Percentage of patients aged 18 years and older seen during the submitting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated	-	X	X	X	X

Measure Title	NQF	Quality Number (Q#)	Measure Description	High Priority	Claims			EHR		Registry	QOPI QCDR (by ASCO and ASTRO)
Falls: Screening for Future Fall Risk	0101	318	Percentage of patients 65 years of age and older who were screened for future fall risk during the measurement period	X	-		X		-		
Opioid Therapy Follow-up Evaluation	N/A	408	All patients 18 and older prescribed opiates for longer than six weeks duration who had a follow-up evaluation conducted at least every three months during Opioid Therapy documented in the medical record	-	-		-		X		X
Documentation of Signed Opioid Treatment Agreement	N/A	412	All patients 18 and older prescribed opiates for longer than six weeks duration who signed an opioid treatment agreement at least once during Opioid Therapy documented in the medical record	-	-		-		X		
Evaluation or Interview for Risk of Opioid Misuse	N/A	414	All patients 18 and older prescribed opiates for longer than six weeks duration evaluated for risk of opioid misuse using a brief validated instrument (eg. Opioid Risk Tool, SOAPP-R) or patient interview documented at least once during Opioid Therapy in the medical record	-	-		-		X		
Proportion of Patients who Died from Cancer with more than One Emergency Department Visit in the Last 30 Days of Life	0211	454	Proportion of patients who died from cancer with more than one emergency room visit in the last 30 days of life	-	-		-		X		
Proportion Admitted to the Intensive Care Unit (ICU) in the Last 30 Days of Life	0213	455	Proportion of patients who died from cancer admitted to the ICU in the last 30 days of life	-	-		-		X		
Proportion Not Admitted To Hospice	0215	456	Proportion of patients who died from cancer not admitted to hospice	-	-		-		X		X
Proportion Admitted to Hospice for less than 3 days	0216	457	Proportion of patients who died from cancer, and admitted to hospice and spent less than 3 days there	-	-		-		X		X
Bone Density Evaluation for Patients with Prostate Cancer and Receiving Androgen Deprivation Therapy	N/A	462	Patients determined as having prostate cancer who are currently starting or undergoing androgen deprivation therapy (ADT), for an anticipated period of 12 months or greater and who receive an initial bone density evaluation. The bone density evaluation must be prior to the start of ADT or within 3 months of the start of ADT	-	-		X		-		X